

No Arbitrage and General Semimartingales

Philip Protter *, Kazuhiro Shimbo

ORIE, Cornell University, Rhodes Hall, Ithaca, NY 14853-3801 USA (e-mail: pep4@cornell.edu and ks266@cornell.edu)

March 1, 2006

Abstract

No free lunch with vanishing risk (NFLVR) is known to be equivalent to the existence of an equivalent martingale measure for the price process semimartingale. We give necessary conditions for such a semimartingale to have the property NFLVR. We also extend Novikov's criterion for the stochastic exponential of a local martingale to be a martingale to the general case; that is, the case where the paths need not be continuous.

Key words Equivalent martingale measure, Girsanov's theorem, No Free Lunch with Vanishing Risk, Semimartingales, Novikov's condition

JFL Classification G10

Mathematical Subject Classification (2000) 60G44, 62P05

1 Introduction

The question of whether the absence of arbitrage is equivalent to the existence of an equivalent measure has now been clarified for some time, in the papers of Delbaen and Schachermayer ([4] and [5]). They showed that one has no arbitrage in the sense of *no free lunch with vanishing risk* if and only if there exists an equivalent probability measure rendering the price process a sigma martingale. (In the continuous case, all sigma martingales are local martingales.) Their conditions, known by its acronym NFLVR, imply also that the price process must be a semimartingale, as a consequence of the Bichteler-Dellacherie theorem. Therefore a natural question arises: which semimartingales actually satisfy NFLVR, and thus can be used to model price processes in arbitrage free models? Partial results in this direction have been obtained by E. Strasser (see [24]) in the continuous case, and also by E. Eberlein and J. Jacod (see [8]) in the case of geometric Lévy processes. In the first half of this paper we consider the general situation and obtain primarily necessary conditions for a semimartingale price process to engender a model without arbitrage. Our primary result in this half is Theorem 2.

When dealing with sufficient conditions, some difficult issues arise: how does one find an equivalent sigma martingale measure? Obvious constructions lead to measures which *a priori* could be sub probability measures, and not true probability measures. The Radon Nikodym densities of these measures can often be constructed as stochastic exponentials of local martingales. A classic tool (in the continuous case) used to verify that the exponential of a local martingale is itself a martingale, and not just a supermartingale, is Novikov's theorem. Often Novikov's theorem is insufficient, but it is always appealing due to its simple nature and ease of computation. In the second half of this paper we propose an analog of Novikov's criterion for the general case (that is, the case with jumps). Our results build on the pioneering work of J. Mémin, A.S. Shiryaev, and their co-authors. Our primary result in this half of the paper is Theorem 6.

* Supported in part by NSF grant DMS-0202958 and NSA grant H98230-06-1-0079

Acknowledgements We wish to thank Jean Jacod for enlightening discussions concerning the first half of this paper. We also wish to thank Darrell Duffie for complaining years ago to the first author that there was no good Novikov type criterion for the general case.

2 Necessary Conditions for No Arbitrage

2.1 The Continuous Case

Let $X_t = X_0 + M_t + A_t$, $t \geq 0$ be a continuous semimartingale on a filtered probability space $(\Omega, \mathcal{F}, \mathbb{F}, P)$ where $\mathbb{F} = (\mathcal{F}_t)_{t \geq 0}$. Here M represents the continuous local martingale part and A is a process with paths of finite variation on compact time sets, almost surely. We seek necessary and sufficient conditions such that there exists an equivalent probability measure P^* such that X is a P^* sigma martingale. Since X is continuous, and since all continuous sigma martingales are in fact local martingales, we need only concern ourselves with local martingales.

Theorem 1 *Let $X_t = X_0 + M_t + A_t$, $0 \leq t \leq T$ be a continuous semimartingale on a filtered probability space $(\Omega, \mathcal{F}, \mathbb{F}, P)$ where $\mathbb{F} = (\mathcal{F}_t)_{0 \leq t \leq T}$. Let $C_t = [X, X]_t = [M, M]_t$, $0 \leq t \leq T$. There exists an equivalent probability measure P^* on \mathcal{F}_T such that X is a P^* sigma martingale only if*

1. $dA \ll dC$ a.s.;
2. If J is such that $A_t = \int_0^t J_s dC_s$ for $0 \leq t \leq T$, then $\int_0^T J_s^2 dC_s < \infty$ a.s.;

If in addition one has the condition below, then we have sufficient conditions for there to exist an equivalent probability measure P^ on \mathcal{F}_T such that X is a P^* sigma martingale:*

3. $E\{\mathcal{E}(-J \cdot M)_T\} = 1$, where $\mathcal{E}(U)$ denotes the stochastic exponential of a semimartingale U .

Proof Suppose there exists P^* equivalent to P such that X is a P^* local martingale. Let $Z = \frac{dP^*}{dP}$ and let $Z_t = E\{Z|\mathcal{F}_t\}$ for all t , $0 \leq t \leq T$. We then have, by Girsanov's theorem, that the decomposition of X under P^* is given by:

$$X_t = X_0 + \left\{ M_t - \int_0^t \frac{1}{Z_s} d[Z, M]_s \right\} + \left\{ A_t + \int_0^t \frac{1}{Z_s} d[Z, M]_s \right\}. \quad (1)$$

Since X is a P^* local martingale and continuous semimartingales have unique decompositions of the type (1), we conclude that we must have

$$A_t = - \int_0^t \frac{1}{Z_s} d[Z, M]_s \quad (2)$$

and since further, by the Kunita-Watanabe inequality, we have $d[Z, M] \ll d[M, M]$ a.s., we conclude that for some predictable process J that

$$A_t = \int_0^t J_s d[M, M]_s. \quad (3)$$

Since Z is a strictly positive P martingale, we can write it as a solution of an exponential equation. (Note that even though X is assumed to be a continuous semimartingale, that does not imply that Z too must be continuous.)

$$Z_t = 1 + \int_0^t Z_{s-} dY_s, \quad Z_0 = 1 \quad (4)$$

for a local martingale Y with $Y_0 = 0$. This Y is often called a stochastic logarithm of Z and is given by $Y = \int_0^t (1/Z_{s-}) dZ_s$. A local martingale Y has a decomposition $Y = Y^c + Y^d$ where Y^c is a continuous martingale part of Y and Y^d is a purely discontinuous martingale part of Y . (See [9, P.85]) Since Y^c is continuous, it is locally square integrable local martingale. Therefore we have a unique representation of the form

$$Y_t^c = \int_0^t H_s dM_s + N_t^c \quad (5)$$

where H is a predictable process such that the stochastic integral in (5) exists, and N^c is a continuous local martingale orthogonal to $H \cdot M$ in the sense that $[H \cdot M, N]^c$ is a local martingale. Since the stochastic integral

in (5) exists, we have of necessity that $\int_0^t H_s^2 d[M, M]_s < \infty$ a.s. for each t , $0 \leq t \leq T$. Let's let $N = N_t^c + Y_t^d$. Then

$$[H \cdot M, N] = [H \cdot M, N^c] + [H \cdot M, Y^d] = [H \cdot M, N^c]. \quad (6)$$

It follows that $[H \cdot M, N]$ is also a local martingale and we have a decomposition of Y into two orthogonal components:

$$Y_t = \left(\int_0^t H_s dM_s + N_t^c \right) + Y_t^d = \int_0^t H_s dM_s + (N_t^c + Y_t^d) = \int_0^t H_s dM_s + N_t \quad (7)$$

We next apply the Meyer-Girsanov theorem to calculate the decomposition of X under P^* . (Since M is continuous there is no issue about the existence of $d\langle Z, M \rangle_s$.) We get:

$$X_t = X_0 + \left\{ M_t - \int_0^t \frac{1}{Z_{s-}} d\langle Z, M \rangle_s \right\} + \left\{ A_t + \int_0^t \frac{1}{Z_{s-}} d\langle Z, M \rangle_s \right\}.$$

By the uniqueness of the decomposition, we must have

$$\begin{aligned} A_t &= - \int_0^t \frac{1}{Z_{s-}} d\langle Z, M \rangle_s = -\langle M, Y \rangle = -\langle M, H \cdot M + N \rangle \\ &= - \int_0^t H_s d[M, M]_s = \int_0^t J_s dC_s, \end{aligned}$$

and by the definition of C we have $-H = J$, $d[M, M]_s dP$ almost everywhere. Thus $\int_0^t J_s^2 d[M, M]_s < \infty$ for all t , $0 \leq t \leq T$. This gives the necessity.

For the sufficiency, let us take Z to be the stochastic exponential of $-J \cdot M$. Applying the Meyer-Girsanov theorem we again have

$$X_t = X_0 + \left\{ M_t - \int_0^t \frac{1}{Z_{s-}} d\langle Z, M \rangle_s \right\} + \left\{ A_t + \int_0^t \frac{1}{Z_{s-}} d\langle Z, M \rangle_s \right\}$$

and by construction we have that $\left\{ A_t + \int_0^t \frac{1}{Z_{s-}} d\langle Z, M \rangle_s \right\} = 0$. The process Z is a strictly positive local martingale with $Z_0 = 1$, hence it is a positive supermartingale, and it is a martingale as soon as $E\{Z_t\} = 1$ for all t , $0 \leq t \leq T$. If Z is known to be a martingale on $[0, T]$ then we define P^* by $dP^* = Z_T dP$, and we can conclude that $\left\{ M_t - \int_0^t \frac{1}{Z_{s-}} d\langle Z, M \rangle_s \right\}$ is a local martingale under P^* . However the third hypothesis guarantees that Z is a martingale and hence that P^* is a probability measure (and not a sub probability measure), and we have sufficiency.

Remark 1 The sufficiency is not as useful in practice as it might seem. The first two conditions should be, in principle, possible to verify, but the third condition is in general not. Depending on the structure of Y , different techniques are available. An obvious one is Novikov's condition, but while easy to state, this too is difficult to verify in practice.

Remark 2 If condition (2) of Theorem 1 is satisfied for **all** ω (instead of P -a.s.), then condition (3) is automatically satisfied. (See, for example, [17]) This is sufficient but not necessary in general. This difference seems subtle but plays an important role. Essentially this is because a probability measure P^* such that X is P^* -local martingale, if it exists, is not necessarily equivalent to P in general a priori.

Remark 3 Condition (1) is often called a structure condition (SC) in the literature. For example, see for example Schweizer [23, page 1538]. Also see Jarrow and Protter (2004) [10] for a constructive example of an arbitrage opportunity when this condition is violated.

Remark 4 In an interesting paper, Strasser [24] discusses a similar problem in the case of continuous semimartingales. She focuses on the condition (1), and does not take the approach we take here.

2.2 General Case

The techniques used in the continuous case break down in the general case (ie, the case with jumps). The reason is that to use formally the same ideas, one would need to use the Meyer-Girsanov theorem, which requires the existence of the process $\langle Z, M \rangle$. When M has continuous paths, such a process always exists, even if Z can have jumps. But if both Z and M have jumps, then the process $\langle Z, M \rangle$ exists if and only if the process $[Z, M]$ is locally integrable, which need not in general be the case. (We mention here that $[Z, M]$ is called locally integrable if there exists a sequence of stopping times $(\tau_n)_{n \geq 1}$ such that $\tau_{n-1} \leq \tau_n$ a.s. for each $n \geq 1$, $\lim_{n \rightarrow \infty} \tau_n \geq T$ a.s., and $E\{[Z, M]_{\tau_n}\} < \infty$ for each $n \geq 1$). A technique developed to circumvent this kind of technical integrability problem is that of random measures, and in particular the use of the *characteristics* of a semimartingale. We assume the reader is familiar with the basic definitions and theorems concerning the characteristics of a semimartingale. We refer the reader to (for example) [9] for an expository treatment of them.

Let X be an arbitrary semimartingale with characteristics (B, C, ν) on our usual filtered probability space $(\Omega, \mathcal{F}, \mathbb{F}, P)$, $\mathbb{F} = (\mathcal{F}_t)_{t \geq 0}$. Then there exists a predictable process A_t with $A_0 = 0$ such that

$$\nu(ds, dx) = K_s(\omega, dx)dA_s(\omega), \quad C_t = \int_0^t c_s dA_s, \quad B_t = \int_0^t b_s dA_s \quad (8)$$

Let P^* be another probability measure equivalent to P . Then of course X is a semimartingale under P^* , with characteristics (B^*, C, ν^*) . (We write C instead of C^* because it is the same process for any equivalent probability measure.) We then know (see Theorem 3.17 on page 170 of [9]) that the random measure ν^* is absolutely continuous with respect to ν , and that there exists a predictable process (predictable in the extended sense) $Y(s, x)_{s \geq 0, x \in \mathbb{R}}$ such that

$$\nu^* = Y \cdot \nu. \quad (9)$$

We have the following theorem, which gives necessary conditions for X to have no arbitrage in the Delbaen-Schachermayer sense of ‘‘No free lunch with vanishing risk,’’ hereafter abbreviated as **NFLVR**. See Delbaen and Schachermayer [4] or alternatively [11].

Theorem 2 *Let X be a P semimartingale with characteristics (B, C, ν) . For X to have an equivalent sigma martingale measure and hence satisfy the NFLVR condition, there must exist a predictable process $\beta = (\beta_t)_{t \geq 0}$ and an (extended) predictable process $Y(\cdot, t, x)$ such that following four conditions are satisfied:*

1. $b_t + \beta_t c_t + \int \{x(Y(t, x) - 1_{\{|x| \leq 1\}})K_t(dx) = 0$; $P(d\omega)dA_s(\omega)$ almost everywhere;
2. $\int_0^T \beta_s^2 dC_s < \infty$, a.s.;
3. $\Delta A_t > 0$ implies that $\int xY(s, x)K(s, dx) = 0$;
4. $\int |x^2| \wedge |x|Y(t, x)K_t(dx) < \infty$, $P(d\omega)dA_s(\omega)$ almost everywhere,

where the predictable process A_t is defined by (8).

Proof Our primary tool will be the Jacod-Mémin version of a Girsanov theorem with characteristics (see Theorem 3.24 on page 172 of [9]). Let P^* be an equivalent sigma martingale measure. Let (B^*, C^*, ν^*) be a characteristics of X under P^* . Then there exist c_s, b_s^*, K_s^* such that

$$C_t^* = c_s dA_s, \quad B_t^* = \int_0^t b_s^* dA_s; \quad \nu^*(ds, dx) = dA_s K_s^*(\omega, dx). \quad (10)$$

Note that in the above, we write c and not c^* , and also A and not A^* , since under our hypothesis, we can take $A^* = A$. In addition the process C does not change under an equivalent change of measure. We next invoke Proposition 6.35 on page 215 of [9] to conclude that X is a P^* sigma martingale if and only if

1. $b_t^* + \int x 1_{\{|x| > 1\}} K_t^*(dx) = 0$, $P(d\omega)dA_s(\omega)$, almost everywhere;
2. When $\Delta A_t > 0$ then $\int x K_t^*(dx) = 0$; and
3. $\int |x^2| \wedge |x| K_t^*(dx) < \infty$, $P(d\omega)dA_s(\omega)$, almost everywhere.

We wish to interpret these three conditions in terms of the original characteristics under P . We know from the continuous case that for C we need a new predictable process coming from the density process Z of $\frac{dP^*}{dP}$, which we denote β , with the property that $\int_0^t \beta_s^2 dC_s = \int_0^t \beta_s^2 c_s dA_s < \infty$ p.s. We also use a key fact that ν^* must be

absolutely continuous with respect to ν and there must exist $Y = Y(s, x)$, predictable in the extended sense, such that

$$\nu^* = Y \cdot \nu. \quad (11)$$

This is proved in Theorem 3.17 on page 170 of [9]. (We remark that both β and Y derive from the P martingale Z where $Z_T = \frac{dP^*}{P}$, with β coming from the continuous martingale part of Z , and Y coming from the ‘purely discontinuous’ part of Z .) Moreover since for any bounded U we have

$$\begin{aligned} \int_0^t \int U(\omega, s, x) \nu^*(dx, ds) &= \int_0^t \int U(\omega, s, x) dA_s K_s^*(dx) \\ &= \int_0^t \int U(\omega, s, x) Y(\omega, s, x) dA_s K_s(dx) \end{aligned}$$

we can conclude that $K^* = Y \cdot K$,

Now we need only to re-express the three conditions in (2.2) to conclude that we must have:

1. $b_t + \beta_t c_t + \int \{x(Y(t, x) - 1_{\{|x| \leq 1\}})\} K_t(dx) = 0$; $P(d\omega) dA_s(\omega)$ almost everywhere;
2. $\int_0^T \beta_s^2 dC_s < \infty$, a.s.;
3. $\Delta A_t > 0$ implies that $\int x K_t^*(dx) = \int x Y(s, x) K(s, dx) = 0$;
4. $\int |x^2| \wedge |x| K_t^*(dx) = \int |x^2| \wedge |x| Y(t, x) K_t(dx) < \infty$, $P(d\omega) dA_s(\omega)$ almost everywhere.

Corollary 1 *Let X be a semimartingale as in Theorem 2. Suppose in addition that \mathbb{F} is a quasi-left continuous filtration. If X is a P^* sigma martingale, then we must have the following three conditions satisfied:*

1. $b_t + \beta_t c_t + \int \{x(Y(t, x) - 1_{\{|x| \leq 1\}})\} K_t(dx) = 0$; $P(d\omega) dA_s(\omega)$ almost everywhere;
2. $\int_0^T \beta_s^2 dC_s < \infty$, a.s.;
3. $\int |x^2| \wedge |x| Y(t, x) K_t(dx) < \infty$, $P(d\omega) dA_s(\omega)$ almost everywhere.

Proof We are able to remove the condition on the jumps of A because if \mathbb{F} is quasi-left continuous, then A does not jump, it being increasing and predictably measurable.

Corollary 2 *Let X be a semimartingale as in Theorem 2. If X is a P^* local martingale, then we must have the following three conditions satisfied:*

1. $b_t + \beta_t c_t + \int \{x(Y(t, x) - 1_{\{|x| \leq 1\}})\} K_t(dx) = 0$; $P(d\omega) dA_s(\omega)$ almost everywhere;
2. $\int_0^T \beta_s^2 dC_s < \infty$, a.s.;
3. $\Delta A_t > 0$ implies that $\int x Y(s, x) K(s, dx) = 0$;

and if the filtration \mathbb{F} is quasi-left continuous, we must have the following two conditions satisfied:

1. $b_t + \beta_t c_t + \int \{x(Y(t, x) - 1_{\{|x| \leq 1\}})\} K_t(dx) = 0$; $P(d\omega) dA_s(\omega)$ almost everywhere;
2. $\int_0^T \beta_s^2 dC_s < \infty$, a.s.;

Proof This follows from Theorem 2 and Proposition 6.35 on page 215 of [9]. That quasi-left continuity of \mathbb{F} implies we can drop the condition on the jumps of A is a trivial consequence of A not having jumps when the filtration is quasi-left continuous.

Remark 5 Comparing Theorem 1 and Theorem 2 illustrates how incompleteness of the market corresponding to the price process X can arise in two different ways. Theorem 1 shows that (in the continuous case) the choice of the orthogonal martingale N is essentially arbitrary, and each such choice potentially leads to a different equivalent probability measure rendering X a local martingale. Theorem 2 shows that in the general case (the case where jumps are present) incompleteness can still arise for the same reasons as in the continuous case, but also because of the jumps, through the choice of Y . Indeed, we are free to change Y appropriately at the cost of changing b . Only if K reduces to a point mass is it then possible to have uniqueness of P^* (and hence market completeness), and then of course only if $C = 0$.

Remark 6 For the special case where X is a geometric Lévy process, Eberlein and Jacod [8] give a necessary and sufficient condition for the existence of an equivalent martingale measure.

We can derive a structure condition for the general case, with an additional hypothesis involving integrability.

Theorem 3 Let X be a special semimartingale with characteristics (B, C, ν) . Then X has a canonical decomposition $X = X_0 + M + A$. Assume $(x^2 \wedge x) * \nu_t < \infty$. If there exists P^* such that X is P^* local martingale, then

$$dA_t \ll d[C_t + (x^2 \wedge x) * \nu_t]. \quad (12)$$

In particular if X is locally square integrable then $\langle M, M \rangle$ exists and

$$dA_t \ll d\langle M, M \rangle_t \quad \text{a.s.} \quad (13)$$

Proof Suppose an equivalent local martingale local measure P^* exists. Let (B, C, ν) and (B^*, C, ν^*) be characteristics of X under P and P^* with truncation function $h(x) = 1_{\{|x| \leq 1\}}$. Let μ be a jump measure of X . Since X is P^* -local martingale, $(x1_{\{|x| > 1\}}) * \mu$ is P^* -locally integrable and X has a representation:

$$X_t = X_0 + X_t^{c*} + x * (\mu - \nu^*)_t \quad (14)$$

Since $P \ll P^*$ by hypothesis, applying Girsanov's theorem (Theorem 3.24 in page 172 of [9]), there exists a predictable process β' and $\mathcal{P} \otimes \mathcal{B}(\mathbb{R})$ measurable non-negative function Y' such that

$$B = B^* + \beta' \cdot [X^c, X^c] + 1_{\{|x| < 1\}}(Y - 1) * d\nu^* \quad (15)$$

$$\nu = Y' \cdot \nu^* \quad (16)$$

where $|1_{\{|x| \geq 1\}}(Y - 1)| * \nu_t^* < \infty$ P^* -a.s. Under P , since we assume X is a special semimartingale, $x1_{\{|x| > 1\}} * \mu$ is P -locally integrable and

$$X_t = X_0 + X_t^c + x * (\mu - \nu)_t + x * (\nu - \nu^*)_t + \int_0^t \beta'_s dC_s = X_0 + M_t + A_t. \quad (17)$$

By the uniqueness of Doob-Meyer decomposition, we have

$$A_t = x(1 - Y') * \nu_t + \int_0^t \beta'_s dC_s \quad (18)$$

Then clearly $A_t \ll d((x^2 \wedge x) * \nu_t + C_t)$. Finally suppose in addition that X is locally square integrable. Then $[M, M]$ is locally integrable and $\langle M, M \rangle = C + x^2 * \nu$ exists. It is clear from (18) that $dA_t \ll d\langle M, M \rangle_t$, a.s.

Remark 7 The case when X is locally bounded (and hence X is a special semimartingale such that M is an automatically locally square integrable local martingale is shown by Delbaen and Schachermayer [6, Theorem 3.5]. Theorem 3 extends their result to the case when X is not necessarily locally bounded. In addition, Theorem 3 does not depend on the notion of admissibility.

Remark 8 The structure condition has a clear economic interpretation. On the set E such that $\int_E d\langle M, M \rangle = 0$, M is constant and $P(\int_E \int_{\mathbb{R}} |x| \mu(dx, dt) = 0) = 1$ where μ is a jump measure of X . Therefore any trading strategy supported on E is risk-free in the sense that any movement of X comes from the predictable component A and hence we can construct a trading strategy which takes advantage of the information of an infinitesimal future. Indeed it is easy to construct such a trading strategy to exploit an arbitrage opportunity if $dA \ll d\langle M, M \rangle$: Consider a price process X on a finite time horizon $[0, T]$. Without loss of generality, we assume that A is an increasing predictable process. Suppose there exists a set $E \in \mathcal{B}(\mathbb{R}_+)$ such that $\int_E d\langle M, M \rangle_s = 0$ but $P(\int_E dA_s > 0) > \eta$ for some $\eta > 0$. Let $A_t^c = A_t - \sum_{0 \leq s \leq t} \Delta A_s$. Let h_t be a predictable process defined by

$$h_t = A_t^c 1_{E \cap \{t: \Delta A_t = 0\}} + \text{sgn}(\Delta A_t) 1_{\{\Delta A_t \neq 0\}}. \quad (19)$$

The following equation is well defined:

$$\int_0^T h_s dX_s = \int_E h_s dM_s + \int_{E \cap \{\Delta A_t = 0\}} A_s^c dA_s^c + \sum_{s \leq T} |\Delta A_s| = \frac{1}{2} \int_E d((A_s^c)^2) + \sum_{s \leq T} |\Delta A_s| \quad (20)$$

Therefore $P(\int_0^T h_s dX_s \geq 0) = 1$ and $P(\int_0^T h_s dX_s > 0) > \eta > 0$. Since $(h \cdot X)_t \geq 0$ for all $t \in [0, T]$, h is a 0-admissible trading strategy and hence this is an arbitrage opportunity.

3 Stochastic exponential of local martingales

3.1 Definition and notations

One of the key components of the sufficient conditions for no arbitrage is that a martingale density Z be a true martingale. However it is not easy to verify this directly in general. The literature is rich on this topic especially for the case when a martingale density Z is continuous. For example Novikov [19], Kazamaki [13], [14], [15], Cherney & Shiryaev [3] studied this question for the continuous case and derived several sufficient conditions in terms of integrability conditions. Mémin [18], Lépingle and Mémin [16], and Kallsen and Shiryaev [12] studied the same question in a general (non-continuous) setting.

The purpose of this section is to show that a formula similar to the famous Novikov condition works in a general setting. More precisely we want to show that a Novikov-type condition $E[\exp\{c\langle M, M \rangle\}] < \infty$ for some c is sufficient to show that $\mathcal{E}(M)$ is a martingale. This condition belongs to the *predictable type* introduced by Revuz and Yor [21].

It should be noted that a Novikov-type condition is often difficult (even the in continuous case) to apply directly. Therefore a common use of this type of condition occurs together with a localization argument. We illustrate this with Example 3.

Let $M = \{M_t\}_{t \geq 0}$ be a càdlàg local martingale vanishing at 0 on a given filtered probability space $(\Omega, \mathcal{F}_t, \mathbb{F}, P)$. A process $X = \{X_t\}_{t \geq 0}$ defined by

$$X_t = \exp \left\{ M_t - \frac{1}{2} [M, M]_t^c \right\} \prod_{s \leq t} (1 + \Delta M_s) \exp(-\Delta M_s) \quad (21)$$

is called a *Doléans exponential* or the stochastic exponential of M , and it is denoted by $\mathcal{E}(M)_t$. X_t is also given as a solution of the stochastic differential equation

$$dX_t = X_{t-} dM_t, \quad X_0 = 1, \quad (22)$$

Since X_- is left continuous and therefore locally bounded, it follows that $\mathcal{E}(M)$ is a local martingale in all cases. When $\Delta M_t > -1$ for all t a.s., it is a positive local martingale. By Fatou's lemma, $\mathcal{E}(M)$ is also a positive supermartingale. Throughout this paper, we assume that $\Delta M_t > -1$.

3.2 Results

3.2.1 Lévy process and additive martingales We start with a basic result shown by Lépingle and Mémin [16].

Theorem 4 (Lépingle and Mémin) *Let M be a local martingale. If the compensator C of the process*

$$\langle M^c, M^c \rangle_t + \sum_{s \leq t} \Delta M_s^2 1_{\{|\Delta M_s| \leq 1\}} + \sum_{s \leq t} \Delta M_s 1_{\{|\Delta M_s| > 1\}} \quad (23)$$

is bounded, then $E[\sqrt{[\mathcal{E}(M), \mathcal{E}(M)]_t}] < \infty$. In particular $\mathcal{E}(M)_t$ is a martingale.

Proof See Lépingle and Mémin [16].

Although the requirement of boundedness looks strong, it is enough to show the following well known fact:

Corollary 3 *Let M be a Lévy martingale. Then $\mathcal{E}(M)$ is a martingale.*

Proof Fix $T > 0$ and let $M_t = M_t^T$. Then by the Lévy decomposition theorem,

$$M_t = W_t + \int_{|x| < 1} x (N(\cdot, [0, t], dx) - t\nu(dx)) + \left(\sum_{0 < s \leq t} \Delta M_s 1_{\{|\Delta M_s| \geq 1\}} - \alpha t \right)$$

where W is a Brownian motion, N is a Poisson random measure with mean measure $dt \nu(dx)$, $\int 1 \wedge x^2 \nu(dx) < \infty$ and $\alpha t = E(\sum_{0 < s \leq t} \Delta M_s 1_{\{|\Delta M_s| \geq 1\}})$. Then the compensator of $[W, W]_t$, $\sum_{0 < u \leq t} |\Delta M_u^1|$, $\sum_{0 < u \leq t} (\Delta M_u^2)^2$ are bounded by T , $T \int_{|x| < 1} x^2 \nu(dx)$ and αT respectively. Therefore the preceding theorem implies that M_t^T is a uniformly integrable martingale for all $T \in \mathbb{R}_+$. Thus M is a martingale.

3.2.2 A general result

Theorem 5 Let M be a martingale such that $\Delta M > -1$. If the process

$$A_t = \frac{1}{2} \langle M^c, M^c \rangle_t + \sum_{s \leq t \wedge T} \{(1 + \Delta M_s) \ln(1 + \Delta M_s) - \Delta M_s\} \quad (24)$$

has compensator $B = (B_t)_{t \geq 0}$ which satisfies

$$E [e^{B_\infty}] < \infty, \quad (25)$$

then $\mathcal{E}(M)$ is a uniformly integrable martingale and $\mathcal{E}(M)_\infty > 0$ a.s.

Proof See Lépingle and Mémin [16, III.1]. Note that $\langle M^c, M^c \rangle_t = [M, M]_t^c$, which is the way it is denoted in [20]. We change the notation here for consistency with Theorem which follows.

For a local martingale M , we let M^d denote its ‘purely discontinuous’ part; that is, $M = M^c + M^d$, where M^c is a local martingale with continuous paths, and $[M^c, M^d]$ is also a (possibly zero) local martingale. (See, e.g., [20, page 193] for the case of a square integrable martingale M , and [7] for the more general case of a local martingale.)

Theorem 6 Let M be a locally square integrable martingale such that $\Delta M > -1$. If

$$E \left[e^{\frac{1}{2} \langle M^c, M^c \rangle_T + \langle M^d, M^d \rangle_T} \right] < \infty, \quad (26)$$

where M^c and M^d are continuous and purely discontinuous martingale parts of M , then $\mathcal{E}(M)$ is martingale on $[0, T]$, where T can be ∞ .

Proof Let $f(x) = (1+x) \ln(1+x) - x$, $g(x) = x^2$. For $x > -1$, $g(x) \geq f(x)$. It follows that

$$L_t \triangleq \sum_{s \leq t} \{g(\Delta M_s) - f(\Delta M_s)\} \geq 0 \quad (27)$$

is a locally integrable increasing process and there exists a compensator \tilde{L}_t of L_t by the Doob-Meyer decomposition. Since $\tilde{L}_t = \sum_{s \leq t} \widetilde{g(\Delta M_s)} - \sum_{s \leq t} \widetilde{f(\Delta M_s)} \geq 0$,

$$\sum_{s \leq t} \widetilde{f(\Delta M_s)} \leq \sum_{s \leq t} \widetilde{g(\Delta M_s)} = \sum_{s \leq t} \widetilde{\Delta[M, M]_s} = \langle M^d, M^d \rangle_t$$

Thus $E \left[e^{\frac{1}{2} \langle M^c, M^c \rangle_T + \langle M^d, M^d \rangle_T} \right] < \infty$ implies the conditions of Theorem 5 and $\mathcal{E}(M)$ is a martingale on $[0, T]$.

A natural question is whether we can improve the constant multiplying $\langle M^d, M^d \rangle$. The following example shows that the answer is negative. Namely $E (e^{(1-\varepsilon)\langle M, M \rangle_\infty}) < \infty$ for any $\varepsilon > 0$ is not sufficient in general.

Example 1 ($\alpha < 1$ is not sufficient) Let $(N_t)_{t \geq 0}$ be a standard Poisson process. Define $T^b = \inf \{s : N_s - (1-b)s = 1\}$ for $b \in (0, 1)$. Let $U_n = \inf_s \{s : N_s = n\}$. Then $P(T^b = \infty) = 0$ since $\{T^b = \infty\} = \bigcap_n \left\{ U_n \geq \frac{n-1}{1-b} \right\}$ and $\frac{U_n}{n} \rightarrow 1$ almost surely. Then

$$N_{T^b} - (1-b)T^b = 1 \quad \text{a.s.} \quad (28)$$

The moment generating function of $N_t - (1-b)t$ exists and for any $\lambda \in \mathbb{R}$

$$E [\exp \{-\lambda [N_t - (1-b)t]\}] = e^{\lambda(1-b)t} E [\exp(-\lambda N_t)] = \exp \{tf(\lambda)\} \quad (29)$$

where

$$f(\lambda) = e^{-\lambda} + \lambda(1-b) - 1 \quad (30)$$

and $Z_t := \exp \{-\lambda [N_t - (1-b)t] - tf(\lambda)\}$ is a martingale. Since Z_t is non-negative, by Doob’s supermartingale inequality, $E(Z_{T^b}) \leq E(Z_0) = 1$. From (28), we obtain

$$E [\exp \{-\lambda [N_{T^b} - (1-b)T^b] - T^b f(\lambda)\}] = E [\exp \{-\lambda - T^b f(\lambda)\}] \leq 1$$

$$E \left[e^{-T^b f(\lambda)} \right] \leq e^\lambda. \quad (31)$$

Now define $M_t = -a(N_t - t)^{T^b}$ where $a \in (0, 1)$. M_t is martingale and $\mathcal{E}(M)_t = \exp\{N_{t \wedge T^b} \ln(1-a) + a(t \wedge T^b)\}$.

$$\begin{aligned} E[\mathcal{E}(M)_{T^b}] &= E[\exp\{N_{T^b} \ln(1-a) + aT^b\}] = E[\exp\{1 + (1-b)T^b\} \ln(1-a) + aT^b] \\ &= (1-a)E[\exp\{T^b\{(1-b)\ln(1-a) + a\}\}] \end{aligned} \quad (32)$$

Let $\lambda^* = \ln\left(\frac{1}{1-a} \frac{1+b}{2}\right)$. Then by (30), $-f(\lambda^*) = -\frac{1-a}{(1+b)/2} + (1-b) \ln\left(\frac{1-a}{(1+b)/2}\right) + 1$. Next define

$$k(a, b) = -f(\lambda^*) - \{(1-b)\ln(1-a) + a\}$$

Simplifying terms,

$$k(a, b) = (1-b) \ln\left(\frac{1}{(1+b)/2}\right) - (1-a) \left(\frac{1}{(1+b)/2} - 1\right). \quad (33)$$

Let $g(b) = 1 - \frac{(1-b)\ln\{2/(1+b)\}}{2/(1+b)-1}$. $k(a, b) > 0$ if $a > g(b)$. Observe that on $\{b : 0 < b < 1\}$, $g(b)$ is an increasing function of b and $1 - \ln 2 < g(b) < 1$. Thus for every $b \in (0, 1)$, there exists $a^* \in (0, 1)$ such that for all $a \geq a^*$, $k(a, b) > 0$. Fix b and choose a so that $k(a, b) > 0$. Then by (31),

$$\begin{aligned} E[\mathcal{E}(M)_{T^b}] &= (1-a)E[\exp\{T^b\{(1-b)\ln(1-a) + a\}\}] \\ &\leq (1-a)E[\exp\{-T^b f(\lambda^*)\}] \\ &\leq (1-a)e^{\lambda^*} = \frac{1+b}{2} < 1 \end{aligned} \quad (34)$$

$\mathcal{E}(M)$ is not a uniformly integrable martingale. $\langle M, M \rangle_t = a^2(t \wedge T_b)$ since a stopped predictable process is still predictable process. Finally, define $h(b)$ by

$$h(b) = b + (1-b)\ln(1-b) = -f(-\ln(1-b)) = \max_\lambda \{-f(\lambda)\}$$

For all $b : 0 < b < 1$, $h(b) < g(b)^2$. However for every $\varepsilon > 0$, there exists $b_\varepsilon^* < 1$ such that $b > b_\varepsilon^*$ implies $h(b) > (1-\varepsilon)g(b)^2$. Fix $\varepsilon > 0$. Let's choose $b' \in (b_\varepsilon^*, 1)$ and $a' \in (g(b'), \sqrt{h(b')/(1-\varepsilon)}) \subset (0, 1)$ so that

$$\begin{cases} a' &> g(b') \\ h(b') &> (1-\varepsilon)a'^2 \end{cases} \quad (35)$$

By (31),

$$E \left[e^{(1-\varepsilon)\langle M, M \rangle_\infty} \right] = E \left[e^{(1-\varepsilon)a'^2 T_{b'}} \right] \leq E \left[e^{(1-\varepsilon)h(b') T_{b'}} \right] < \frac{1}{1-b'} < \infty \quad (36)$$

As seen in this example, jumps with size close to -1 prohibit the improvement of the condition. The following corollary shows that if ΔM is bounded away from -1 , we can improve the results with a constant α . In particular, if $\Delta M > 0$, then $\alpha = 1/2$.

Corollary 4 Fix $\varepsilon \in [0, 1]$. Let M be a locally square integrable martingale such that $\Delta M > -1 + \varepsilon$. Then there exists $\alpha(\varepsilon) \in [1/2, 1]$ such that

$$E \left[e^{\frac{1}{2}\langle M^c, M^c \rangle_T + \alpha(\varepsilon)\langle M^d, M^d \rangle_T} \right] < \infty, \quad (37)$$

implies that $\mathcal{E}(M)_t$ is martingale on $[0, T]$, where T can be ∞ .

Proof Let $f(x) = (1+x)\ln(1+x) - x$, Then there exists $\alpha(\varepsilon) = \inf\{a : ax^2 - f(x) \geq 0 \text{ on } x > -1 + \varepsilon\}$ such that $\alpha(\varepsilon) \in [1/2, 1]$. Especially when $\varepsilon = 1$, we can take $\alpha(\varepsilon) = 1/2$ and $\alpha(\varepsilon)$ is a decreasing function of ε . Let $g(x) = \alpha(\varepsilon)x^2$ For $x > 0$, $g(x) \geq f(x)$. It follows that

$$L_t \triangleq \sum_{s \leq t} \{g(\Delta M_s) - f(\Delta M_s)\} \geq 0$$

is a locally integrable increasing process and there exists a compensator \tilde{L}_t of L_t by the Doob-Meyer decomposition. Since $\tilde{L}_t = \sum_{s \leq t} \widetilde{g(\Delta M_s)} - \sum_{s \leq t} \widetilde{f(\Delta M_s)} \geq 0$

$$\sum_{s \leq t} \widetilde{f(\Delta M_s)} \leq \sum_{s \leq t} \widetilde{g(\Delta M_s)} = \sum_{s \leq t} \widetilde{\Delta[M, M]_s} = \langle M^d, M^d \rangle_t$$

Thus $E \left[\exp\{1/2 \langle M^c, M^c \rangle_t T + \alpha(\varepsilon) \langle M^d, M^d \rangle_T\} \right] < \infty$ implies the condition of theorem 5 and hence $\mathcal{E}(M)$ is martingale on $[0, T]$.

Remark 9 This integrability approach provides sufficient but not necessary conditions. While it is possible to derive a sequence of sufficient conditions converging in some sense to a necessary and sufficient condition, those stronger conditions become more difficult to verify at the same time. For details on this issue, see Kallsen and Shiryaev [12].

In the continuous framework, it is well known that the Novikov condition is not optimal. The symmetric nature of a quadratic variation processes requires that if a continuous martingale M satisfies Novikov's condition, $-M$ has to satisfy Novikov's condition as well. This implies that Novikov's condition is not applicable to identify a class of martingales M such that $\mathcal{E}(M)$ is a martingale but $\mathcal{E}(-M)$ is not. More generally, if there exists a predictable process h_s such that $\int_s h_s dM_s$ is a continuous local martingale satisfying Novikov's condition, then for all integrable predictable $g_s \in L(M)$ such that $\int |g_s|^2 d[M, M]_s = \int h_s^2 d[M, M]_s$, $\mathcal{E}(\int g_s dM_s)$ is a uniformly integrable martingale. See, for example, Stroock [25]. (The authors thank Marc Yor for calling this reference to their attention.)

Some of these examples can be dealt with using a stronger condition derived in an integrability approach, such as Kazamaki's condition. But other examples requires totally different approaches. See for example Lipster and Shiryaev [17], Cheridito, Filipovic, and Yor [2].

Despite these examples showing its limitations, a Novikov-type condition is the kind of condition that we could hope to verify in a practical setting. This is due to the fact that the condition is given in terms of an increasing process and the quadratic variation of $\mathcal{L}og(Z)$, where $\mathcal{L}og(\cdot)$ denotes a stochastic logarithm.

3.3 Examples and applications

The following example shows that when the stochastic exponential comes from a driving Lévy local martingale, then the condition in Theorem 6 becomes easier to compute.

Example 2 Let M be a Lévy local martingale with Lévy triplet (B, C, ν) . Let $h \in L(M)$ be a predictable process such that $X = \int h_s dM_s$ is locally square integrable and $\Delta X_t = h_t \Delta M_t > -1$. Since

$$[X^c, X^c]_t = \int_0^t h_s^2 C ds \quad \langle X^d, X^d \rangle = \int_0^t h_s^2 d\langle M^d, M^d \rangle_s = \int_0^t h_s^2 \left(\int_{\mathbb{R}} x^2 d\nu(dx) \right) ds \quad (38)$$

Let $K = \int_{\mathbb{R}} x^2 \nu(dx)$. Then

$$E \left[\exp \left\{ \left(\frac{1}{2} C + K \right) \int_0^T h_s^2 ds \right\} \right] < \infty \quad (39)$$

then $\mathcal{E}(X)_t$ is a martingale. In particular, if $K < \infty$ and $|h_t|$ is uniformly bounded, $\mathcal{E}(X)_t$ is a martingale.

The following auxiliary lemma is useful in some situations. It allows us to relax the constants that appear in the exponent (e.g. 1 in (26), $\alpha(\varepsilon)$ in Corollary 4 and $1/2 C+K$ in (39) etc).

Lemma 1 *Let M be a local martingale. If there exists $k \in \mathbb{N}_+$ and a partition of $[0, T]$, $0 = \tau_0 < \tau_1 < \dots < \tau_k = T$ where each τ_i is a stopping time such that*

$$E \left[\exp \left[\langle M, M \rangle_{\tau_{m+1}} - \langle M, M \rangle_{\tau_m} \right] \right] < \infty, \quad \forall 1 \leq m \leq k-1, \quad (40)$$

then, $\mathcal{E}(M)$ is a martingale.

Proof Fix m and let $M'_t = M_{t \wedge \tau_{m+1}} - M_{t \wedge \tau_m}$. That is, $M'_t = 0$ if $t < \tau_m$, $M'_t = M_t - M_{\tau_m}$ if $t \in [\tau_m, \tau_{m+1})$, and $M'_t = M_{\tau_{m+1}}$. Since M is a local martingale, M' is also a local martingale. By hypothesis and the previous lemma, $\mathcal{E}(M')_t$ is a martingale.

$$E[\mathcal{E}(M')_t] = 1, \quad \forall t \in [0, t] \quad (41)$$

Let's rewrite M' by $M^{(m)}$. Then for $0 \leq m \leq k-2$

$$\mathcal{E}\left(M^{(m)}\right) \mathcal{E}\left(M^{(m+1)}\right) = \mathcal{E}\left(M^{(m)} + M^{(m+1)} - \left[M^{(m)}, M^{(m+1)}\right]\right) = \mathcal{E}\left(M^{(m)} + M^{(m+1)}\right), \quad (42)$$

and $\mathcal{E}(M_T) = \mathcal{E}\left(\sum_{n=0}^{k-1} M^{(n)}\right)_T$.

$$E[\mathcal{E}(M)_T] = E[E[\mathcal{E}(M)_T | \mathcal{F}_{\tau_{k-1}}]] = E\left[\sum_{n=0}^{k-2} M_t^{(n)} E\left[M_t^{(k-1)} | \mathcal{F}_{\tau_{k-1}}\right]\right] = E[\mathcal{E}(M_{\tau_{k-1}})] \quad (43)$$

Then by induction, $E[\mathcal{E}(M)_T] = E[\mathcal{E}(M)_0] = 1$. Since $\mathcal{E}(M)$ is a local martingale, this implies that it is a martingale.

We can apply Lemma 1 to refine Example 2.

Example 3 In Example 2, if there exists an increasing sequence of stopping times $0 = \tau_0 < \tau_1 \cdots < \tau_k = T$ such that

$$E\left[\exp\left\{\left(\frac{1}{2}C + K\right) \int_{\tau_i}^{\tau_{i+1}} h_s^2 ds\right\}\right] < \infty, \forall i \in \{1, \dots, k-1\} \quad (44)$$

then $\mathcal{E}(X)$ is a martingale on $[0, T]$.

Another application of Lemma 1 yields an extension of Beneš theorem [1]:

Theorem 7 Let $Z_t = \mathcal{E}(h \cdot M)_t$ where M is a Lévy-martingale, $h \in L(M)$ be a predictable adapted process such that $h \cdot M$ is properly defined and $Z_t \geq 0$ for all $t \in [0, T]$. Suppose there exists a $a > 0$ such that $E[e^{a|M_T|}] < \infty$. If there exists a constant K such that

$$|h_t(\omega)| \leq K(1 + \sup_{u \leq t} |M_u(\omega)|^{1/2}) \quad \text{a.s.} \quad (45)$$

for all $t \in [0, T]$ then $\mathcal{E}(h \cdot M)$ is a martingale on $[0, T]$.

Proof Let $\alpha = \sup_{u \leq T} |M_u|^{1/2}$. Then

$$\begin{aligned} \langle h \cdot M, h \cdot M \rangle_{t_{m+1}} - \langle h \cdot M, h \cdot M \rangle_{t_m} &= \int_{t_m}^{t_{m+1}} h_u^2 d\langle M, M \rangle_u = c \int_{t_m}^{t_{m+1}} h_u^2 du \\ &= cK^2(t_{m+1} - t_m)(1 + \alpha)^2 \leq 2cK^2(t_{m+1} - t_m)(1 + \alpha^2) \end{aligned} \quad (46)$$

Therefore

$$E\left[\exp\left(\langle h \cdot M, h \cdot M \rangle_{t_{m+1}} - \langle h \cdot M, h \cdot M \rangle_{t_m}\right)\right] \leq c_1 E\left[\exp(c_2 \alpha^2)\right] \quad (47)$$

where $c_1 = cK^2$, $c_2 = 2cK^2(t_{m+1} - t_m)$. By the hypothesis that $e^{a|M_t|} \in L_1$ for some $a \in \mathbb{R}_+$ and Jensen's inequality, $e^{a|M_t|}$ is a submartingale. By Doob's maximal inequality,

$$E\left[\exp(c_2 \alpha^2)\right] \leq 4E\left[e^{c_2 |M_T|}\right] < \infty, \quad (48)$$

since we can take c_2 arbitrarily small by fining a partition. Therefore by the previous lemma, $\mathcal{E}(h \cdot M)$ is a martingale.

Remark 10 The original Beneš theorem requires $e^{a(M_T)^2} \in L_1$ for some a , instead of $e^{a|M_T|} \in L_1$, while it assumes further that

$$|h_t(\omega)| \leq K(1 + \sup_{u \leq t} |M_u(\omega)|) \quad \text{a.s.} \quad (49)$$

is a sufficient condition for h (note that there is no square root in the original condition). With a proof similar in spirit to Theorem 7, we can show that this still holds even if M is not continuous. However the condition $e^{a|M_t|} \in L_1$ is satisfied only when $\nu(dx) = 0$, which of course implies that M is a Brownian motion as Lemma 2 indicates.

The following lemma is well known, but hard to find explicitly stated.

Lemma 2 Let X be a Lévy process such that its Lévy measure $\nu(dx) \not\equiv 0$. If $p > 1$, then $\exp[a|X_t|^p] \notin L_1$ for all $t > 0, a > 0$.

Proof Suppose there exists ϵ such that $\nu([\epsilon, \infty)) \not\equiv 0$. Let $X_t^{(2)} = \sum_{s \leq t} 1_{[\epsilon, \infty)}(\Delta X_s)$ and $X_t^{(1)} = X_t - X_t^{(2)}$. Then $X^{(1)}$ and $X^{(2)}$ are independent. $P(X_t^{(1)} > 0) > 0$. If $P(X_t^{(2)} = 0) = 1$ then

$$E[\exp\{a|X_t|^p\}] = E[\exp\{a|X_t^{(1)}|^p\}] \geq E[\exp\{a|\epsilon\mu([\epsilon, \infty))|^p\}] = \infty \quad (50)$$

Consider the case $P(X_t^{(2)} = 0) \neq 1$. For $a, b \in \mathbb{R}$, $|a + b|^p \geq |a|1_{\{a, b < 0\}} \cup \{a, b > 0\}$. Therefore

$$\begin{aligned} E[\exp\{a|X_t|^p\}] &\geq E[\exp\{a|X_t^{(1)}|^p\} 1_{\{X_t^{(1)}, X_t^{(2)} < 0\} \cup \{X_t^{(1)}, X_t^{(2)} > 0\}}] \\ &\geq \min[P(X_t^{(2)} > 0), P(X_t^{(2)} < 0)] E[\exp\{a|X_t^{(1)}|^p\}] \\ &\geq \min[P(X_t^{(2)} > 0), P(X_t^{(2)} < 0)] E[\exp\{a|\epsilon\mu_T([\epsilon, \infty))\}|^p] \end{aligned} \quad (51)$$

The case that X has jumps with negative size only is similar.

Lemma 3 Let $Z_t = \mathcal{E}(h \cdot M)_t$ where M is a Lévy-martingale, $h \in L(M)$ is a predictable process such that $h \cdot M$ is well defined and $Z_t \geq 0$. Suppose there exists constants K and p such that

$$|h_s(\omega)| \leq K(1 + ([M, M]_T)^p), \quad (52)$$

and $E[e^{a[M, M]_T^{2p}}] < \infty$ for some $a \in \mathbb{R}_+$, almost surely for all $s \in [0, T]$. Then $\mathcal{E}(M)$ is a martingale on $[0, T]$.

Proof Let $\alpha = [M, M]_T^p$. Then

$$\begin{aligned} \langle h \cdot M, h \cdot M \rangle_{t_{m+1}} - \langle h \cdot M, h \cdot M \rangle_{t_m} &= \int_{t_m}^{t_{m+1}} h_u^2 d\langle M, M \rangle_u = c \int_{t_m}^{t_{m+1}} h_u^2 du \\ &= cK^2(t_{m+1} - t_m)(1 + \alpha)^2 \leq 2cK^2(t_{m+1} - t_m)(1 + \alpha^2) \end{aligned} \quad (53)$$

then the hypothesis that

$$E[e^{c_2[M, M]_T^{2p}}] < \infty \quad (54)$$

implies the desired result.

Example 4 Let M be a compensated compound Poisson process and let h be a predictable process satisfying

$$|h_t(\omega)| \leq K(1 + \sum_{u \leq t} \Delta |M_u|^2). \quad (55)$$

Since $[M, M]_T = \sum_{s \leq T} (\Delta M_s)^2$,

$$E[\exp\{c[M, M]_T\}] = E\left[E\left(\exp\left\{c \sum_{j=1}^{N_t} \xi_j^2\right\} \middle| \sigma(N_t)\right)\right] = E\left[\left(E\left(e^{c\xi_1^2}\right)\right)^{N_t}\right] < \infty, \quad (56)$$

provided $e^{c\xi_1^2} \in L_1$ for some $c > 0$. For example if ξ is bounded or $\xi = |\chi|$ where χ is normally distributed, then this condition is satisfied. In this case, $\mathcal{E}(h \cdot M)$ is a martingale.

References

1. Beneš, V. E. “Existence of optimal stochastic control laws,” *SIAM J. Control* 9, 1971, 446-472.
2. P. Cheridito, D. Filipovic, and M. Yor, “Equivalent and absolutely continuous measure changes for jump-diffusion processes,” *Ann. Applied Probability* 15, 2005, 1713–1732.
3. A.S. Cherney and A. N. Shiryaev, “On Criteria for the Uniform Integrability of Brownian Stochastic Exponentials,” *Optimal Control and Partial Differential Equations*, IOS Press. In honour of Alain Bensoussan’s 60th birthday, 2001, 80-92.
4. F. Delbaen and W. Schachermayer, “A General Version of the Fundamental Theorem of Asset Pricing,” *Math. Ann.* 300, 1994, 463–520.
5. F. Delbaen and W. Schachermayer, “The Fundamental Theorem for Unbounded Stochastic Processes,” *Math. Ann.* 312, 1998, 215–250.
6. F. Delbaen and W. Schachermayer, “The Existence of Absolutely Continuous Local Martingale Measures,” *Ann. Applied Probability* 5, 1995, 926–945.
7. C. Dellacherie and P.A. Meyer, *Probabilities and Potential B*, Volume 72 of *North-Holland Mathematics Studies*, North-Holland, Amsterdam, 1978.
8. Ernst Eberlein and Jean Jacod, “On the ranges of options prices,” *Finance Stochast.* 1, 1997, 131–140.
9. J. Jacod and A.N. Shiryaev, *Limit Theorems for Stochastic Processes, Second Edition*, Springer–Verlag, Heidelberg, 2002.
10. R. Jarrow and P. Protter, “Large Traders, Hidden Arbitrage and Complete Markets,” *Journal of Banking and Finance*, 29: 2803-2820, 2005.
11. R. Jarrow and P. Protter, “An Introduction to Financial Asset Pricing,” to appear in the *Elsevier Handbook of Financial Engineering*.
12. Jan Kallsen and Albert N. Shiryaev, “The cumulant process and Esscher’s change of measure,” *Finance Stoch.*, 6(4):397–428, 2002.
13. N. Kazamaki, “On a problem of Girsanov,” *Tôhoku Math. J.*, 29(4):597–600, 1977.
14. N. Kazamaki, Correction: “On a problem of Girsanov”, (*Tôhoku Math. J. (2)* 29 (1977), no. 4, 597–600), *Tôhoku Math. J. (2)*, 30(1):175, 1978.
15. Norihiko Kazamaki, *Continuous exponential martingales and BMO*, volume 1579 of *Lecture Notes in Mathematics*, Springer-Verlag, Berlin, 1994.
16. Dominique Lépingle and Jean Mémin, “Sur l’intégrabilité uniforme des martingales exponentielles”, *Z. Wahrsch. Verw. Gebiete*, 42(3):175–203, 1978.
17. Robert S. Liptser and Albert N. Shiryaev, *Statistics of random processes. I*, volume 5 of *Applications of Mathematics (New York)*. Springer-Verlag, Berlin, expanded edition, 2001. General theory, Translated from the 1974 Russian original by A. B. Aries, *Stochastic Modeling and Applied Probability*.
18. Jean Mémin, “Décompositions multiplicatives de semimartingales exponentielles et applications”, In *Séminaire de Probabilités, XII (Univ. Strasbourg, Strasbourg, 1976/1977)*, volume 649 of *Lecture Notes in Math.*, pages 35–46. Springer, Berlin, 1978.
19. A. A. Novikov, “On conditions for uniform integrability for continuous exponential martingales.”, In *Stochastic differential systems (Proc. IFIP-WG 7/1 Working Conf., Vilnius, 1978)*, volume 25 of *Lecture Notes in Control and Information Sci.*, pages 304–310. Springer, Berlin, 1980.
20. P. Protter *Stochastic Integration and Differential Equations, Second Edition, Version 2.1*, Springer–Verlag, Heidelberg, 2005.
21. Daniel Revuz and Marc Yor, *Continuous martingales and Brownian motion*, volume 293 of *Grundlehren der Mathematischen Wissenschaften*, Springer-Verlag, Berlin, third edition, 1999.
22. Hiroshi Sato, “Uniform integrability of an additive martingale and its exponential,” *Stochastics Stochastics Rep.*, 30, 1990, 163–169.
23. Martin Schweizer, “Approximating Random Variables by Stochastic Integrals,” *Ann. Probability* 22, 1994, 1536–1575.
24. Eva Strasser, “Characterization of arbitrage-free markets”, *Ann. Appl. Probability* 15, 2005, 116–124.
25. Daniel W. Stroock, *Markov processes from K. Itô’s perspective*, volume 155 of *Annals of Mathematics Studies.*, Princeton University Press, Princeton, NJ, 2003.